For Research Use Only

ZNF618 Polyclonal antibody

Catalog Number:31972-1-AP



Basic Information

 Catalog Number:
 GenBank Accession Number:

 31972-1-AP
 NM_001318042.1

 Size:
 GeneID (NCBI):

 500 ug/ml
 114991

 Source:
 UNIPROT ID:

 Rabbit
 Q5T7WO

IgG zinc finger protein 618
Immunogen Catalog Number: Calculated MW:
AG36487 105kDa, 954aa

Full Name:

Purification Method: Antigen affinity Purification Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500 IF/ICC 1:400-1:1600

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Species Specificity: human

Isotype:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 cancer tissue IF/ICC : A549 cells,

Positive Controls:

WB: MCF-7 cells, SH-SY5Y cells, Caki-2 cells

IHC: human cervical cancer tissue, human stomach

Background Information

Zinc finger protein 618 (ZNF618) is a key protein associated with DNA methylation and acts as a specific 5hmC reader in vivo to regulate UHRF2 function by promoting UHRF2 chromatin localization. In addition, ZNF618 is not only associated with the induction, maintenance or progression of female breast cancer, but is also up-regulated in lung cancer cells.

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 31972-1-AP (ZNF618 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 31972-1-AP (ZNF618 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 31972-1-AP (ZNF618 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed A549 cells using ZNF618 antibody (31972-1-AP) at dilution of 1:800 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-Phalloidin (red).