## For Research Use Only

## CASK Polyclonal antibody

Catalog Number:55277-1-AP

**Featured Product** 



**Basic Information** 

Catalog Number: 55277-1-AP

Size: 600 µg/ml Source: Rabbit Isotype:

otype: Full Name:

G calcium/calmodulin-dependent
serine protein kinase (MAGUK fa

serine protein kinase (MAGUK family) Calculated MW: 105 kDa

Observed MW: 100-110 kDa, 50-55 kDa

GenBank Accession Number:

NM 003688

**UNIPROT ID:** 

014936

GeneID (NCBI):

**Applications** 

Tested Applications: IHC, WB, ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: mouse brain tissue, mouse liver tissue, A549 cells, rat brain tissue

**Purification Method:** 

WB 1:2000-1:10000 IHC 1:500-1:2000

Antigen affinity purification

Recommended Dilutions:

IHC: rat small intestine tissue, mouse kidney tissue, human thyroid cancer tissue

## **Background Information**

CASK, also named as LIN2, belongs to the MAGUK family. It is a multidomain scaffolding protein with a role in synaptic transmembrane protein anchoring and ion channel trafficking. It contributes to neural development and regulation of gene expression via interaction with the transcription factor TRB1. CASK binds to cell-surface proteins, including amyloid precursor protein, neurexins and syndecans. It may mediate a link between the extracellular matrix and the actin cytoskeleton via its interaction with syndecan and with the actin/spectrin-binding protein 4.1. Defects in CASK are the cause of mental retardation X-linked CASK-related (MRXCASK). Defects in CASK are the cause of FG syndrome type 4 (FGS4). This antibody can recognize all the isoform of CASK.

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 55277-1-AP (CASK antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of CASK antibody (55277-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-CASK transfected A549 cells.



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 55277-1-AP (CASK antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human thyroid cancer tissue slide using 55277-1-AP (CASK antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat small intestine tissue slide using 55277-1-AP (CASK antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).