For Research Use Only

HDGF Monoclonal antibody

Catalog Number:60064-1-lg 7 Publications



Basic Information

Catalog Number: 60064-1-lg

Size: 1100 µg/ml Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG1893

GenBank Accession Number:

BC018991 GeneID (NCBI): 3068 **UNIPROT ID:**

Full Name: hepatoma-derived growth factor (high-mobility group protein 1-like)

Calculated MW: 26 kDa

Observed MW: 40 kDa

P51858

Applications

Tested Applications: IHC, WB,ELISA Cited Applications: WB, IF, IHC, CoIP Species Specificity: human, mouse, rat

Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: SMMC-7721 cells, A549 cells, HeLa cells, HepG2 cells, LO2 cells, rat liver tissue

Purification Method:

Protein G purification

Recommended Dilutions:

WB 1:1000-1:5000 IHC 1:50-1:500

CloneNo.:

5A7G3

IHC: human liver cancer tissue, human colon cancer tissue, human kidney tissue

Background Information

HDGF, also named as HMG1L2, belongs to the HDGF family. It is a heparin-binding protein, with mitogenic activity for fibroblasts. HDGF acts as a transcriptional repressor. It plays important roles in the development and progression of cancers. HDGF may have a synergistic effect with VEGF during cancer development and Progression. (PMID:20848225)

Notable Publications

Author	Pubmed ID	Journal	Application
Li San-Zhong SZ	23771798	Med Oncol	IHC
Xin Chen	31358969	Nat Cell Biol	WB
Ye-Wei Zhang	34368121	Front Cell Dev Biol	WB,IF,CoIP

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



SMMC-7721 cells were subjected to SDS PAGE followed by western blot with 60064-1-1g (HDGF Antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 60064-1-1g (HDGF Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 60064-1-1g (HDGF Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).