For Research Use Only

CD14 Monoclonal antibody

Catalog Number: 60253-1-lg 13 Publications



Basic Information

Catalog Number: 60253-1-lg Size:

1000 µ g/ml
Source:

Isotype: IgG1

Immunogen Catalog Number:

AG10693

Mouse

P08571 Full Name: CD14 molecule Calculated MW: 375 aa, 40 kDa

BC010507

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000170458

UNIPROT ID:

Observed MW: 55 kDa

GenBank Accession Number:

Purification Method: Protein G purification

CloneNo.: 2C1D9

Recommended Dilutions: WB 1:500-1:2000 IHC 1:1000-1:4000 IF 1:50-1:500

Applications

Tested Applications:

FC, IF/ICC,IF-P, IHC, WB, ELISA

Cited Applications:

Cell treatment, FC, IF, IHC, WB

Species Specificity: human, mouse Cited Species:

human, mouse, rabbit

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: THP-1 cells, RAW 264.7 cells

IHC: human tonsillitis tissue,

IF: RAW 264.7 cells, human tonsillitis tissue

Background Information

CD14 is a 50-55 kDa glycosylphosphatidylinositol-anchored glycoprotein preferentially expressed on monocytes and macrophages, and at lower levels on granulocytes (PMID: 3385210; 2462937; 7685797). CD14 can also exist as a soluble protein. CD14 acts as a co-receptor for bacterial liposaccharides (LPS) (PMID: 1698311). It plays a major role in the inflammatory response of monocytes to LPS.

Notable Publications

Author	Pubmed ID	Journal	Application
Zeqing Zhai	34480835	Arthritis Rheumatol	IF
Haochen Hui	36208807	Int J Biol Macromol	WB
Jia Xiong	35320754	Mol Cell	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




THP-1 cells were subjected to SDS PAGE followed by western blot with 60253-1-Ig (CD14 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 burs.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60253-1-1g (CD14 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60253-1-1g (CD14 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of RAW 264.7 cells using 60253-1-ig (CD14 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).



1X10^6 THP-1 cells were surface stained with 0.4 ug Anti-Human CD14 (60253-1-Ig, Clone:2C1D9) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were not fixed.