### For Research Use Only

# IL-2 Monoclonal antibody

Catalog Number:60306-1-lg 8 Publications



**Basic Information** 

Catalog Number: 60306-1-lg

Size:
2600 µg/ml
Source:
Mouse
Isotype:
IgG2a

Immunogen Catalog Number: Calculated MW: AG10199 153 aa, 18 kDa

Observed MW: 18 kDa Purification Method:

Protein A purification

CloneNo.: 2F11C5

Recommended Dilutions: WB 1:1000-1:4000 IHC 1:20-1:200 IF 1:200-1:800

**Applications** 

Tested Applications: IF-P, IHC, WB, ELISA Cited Applications:

IHC, WB

Species Specificity:

human
Cited Species:
human, mouse, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, MOLT-4 cells, human peripheral blood leukocyte

IHC : human tonsillitis tissue, human breast cancer

tissue

GenBank Accession Number:

BC070338

3558

P60568

GeneID (NCBI):

UNIPROT ID:

Full Name:

interleukin 2

IF: human tonsillitis tissue,

## **Background Information**

IL2 is a secreted cytokine involved in the operation and regulation of immune system. IL2 plays a pivotal role in T-cell activation and effector functions, including T cell proliferation, interferon-  $\gamma$  production and cytotoxicity. IL2 can stimulate the propagation of lymphocytes and induce the cytotoxic T lymphocytes and lymphokine-activated killer cells for tumor cells. IL2 also influences memory T cell homeostasis through the regulation of memory T cell numbers and enhances major histocompatibility complex II (MHC II) expression on tumor cells.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Chengfen Wang	25264893	PLoS One	WB
Hai long Yang	34657991	Inflammation	WB
Yinrui Guo	34045460	Transl Psychiatry	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 60306-1-1g (IL-2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60306-1-lg (IL2 Antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieved with Tris-EDTA buffer, pH9.0.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60306-1-lg (IL2 Antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieved with Tris-EDTA buffer, pH9.0.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using IL-2 antibody (60306-1-lg, Clone: 2F11C5) at dilution of 1:400 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using IL-2 antibody (60306-1-lg, Clone: 2F11C5) at dilution of 1:400 and CoraLite®488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).