

For Research Use Only

TP63 Monoclonal antibody

Catalog Number: 60332-2-Ig **1 Publications**



Basic Information

Catalog Number: 60332-2-Ig	GenBank Accession Number: BC039815	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 8626	CloneNo.: 4D6G12
Source: Mouse	UNIPROT ID: Q9H3D4	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:2000-1:8000
Isotype: IgG2a	Full Name: tumor protein p63	
Immunogen Catalog Number: AG2791	Calculated MW: 680 aa, 77 kDa	
	Observed MW: 70 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : HaCaT cells, A431 cells, U-937 cells
Cited Applications: WB	IHC : rat skin tissue, human prostate cancer tissue, mouse skin tissue
Species Specificity: human	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

TP63, also named KET, P63, P73H, P73L and TP73L, belongs to the p53 family. It is a homologue of the tumor suppressor p53 and p73 genes. It is involved in malignancy acquisition and maintenance of cells. Unlike p53, the p63 gene encodes multiple isotypes with remarkably divergent abilities to transactivate p53 reporter genes and induce apoptosis. TP63 acts as a sequence specific DNA binding transcriptional activator or repressor. TP63 has 12 isoforms with MW 40kd(P40), 50kd(P60), 63kd(P63) and 73kd(P73L). It is Nuclear stain.

Notable Publications

Author	Pubmed ID	Journal	Application
Yongjie Chen	37654726	Dose Response	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HaCaT cells were subjected to SDS PAGE followed by western blot with 60332-2-Ig (TP63 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded rat skin tissue slide using 60332-2-Ig (TP63 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




A431 cells were subjected to SDS PAGE followed by western blot with 60332-2-Ig (TP63 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse skin tissue slide using 60332-2-Ig (TP63 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



U-937 cells were subjected to SDS PAGE followed by western blot with 60332-2-Ig (TP63 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 60332-2-Ig (TP63 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).