

For Research Use Only

# Cleaved PARP1 Monoclonal antibody, PBS Only (Capture)

Catalog Number: 60555-1-PBS



## Basic Information

<b>Catalog Number:</b> 60555-1-PBS	<b>GenBank Accession Number:</b> BC037545	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 1 mg/ml	<b>GeneID (NCBI):</b> 142	<b>CloneNo.:</b> 4G4C8
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> P09874	
<b>Isotype:</b> IgG1	<b>Full Name:</b> poly (ADP-ribose) polymerase 1	
	<b>Calculated MW:</b> 1014 aa, 113 kDa	
	<b>Observed MW:</b> 89 kDa	

## Applications

**Tested Applications:**  
WB, IHC, IF/ICC, FC (Intra), Indirect ELISA, Sample test

**Species Specificity:**  
human, mouse, rat

## Background Information

PARP1 (poly(ADP-ribose) polymerase 1) is a nuclear enzyme catalyzing the poly(ADP-ribosyl)ation of many key proteins in vivo. The normal function of PARP1 is the routine repair of DNA damage. Activated by DNA strand breaks, the PARP1 is cleaved into an 85 to 89-kDa COOH-terminal fragment and a 24 kDa NH2-terminal peptide by caspases during the apoptotic process. The appearance of PARP fragments is commonly considered an important biomarker of apoptosis. In addition to caspases, other proteases like calpains, cathepsins, granzymes, and matrix metalloproteinases (MMPs) have also been reported to cleave PARP1 and give rise to fragments ranging from 42-89 kDa.

This antibody only recognizes the cleaved form of PARP1 but not full-length PARP1.

## Storage

**Storage:**  
Store at -80°C.  
**The product is shipped with ice packs. Upon receipt, store it immediately at -80°C**

**Storage Buffer:**  
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


# Selected Validation Data




Staurosporine treated and untreated A2780 cells were subjected to SDS PAGE followed by western blot with 60555-1-Ig (Cleaved PARP1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with HRP-conjugated Lamin B1 (HRP-66095) antibody as a loading control. This data was developed using the same antibody clone with 60555-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded Jurkat (left) and Staurosporine treated Jurkat (right) cells slide using 60555-1-Ig (Cleaved PARP1 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60555-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed untreated and 1 μM Staurosporine (3 hours) treated HSC-T6 cells using Cleaved PARP1 antibody (60555-1-Ig, Clone: 4G4C8) at dilution of 1:1000 and Multi-rAb Coralite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004). This data was developed using the same antibody clone with 60555-1-PBS in a different storage buffer formulation.




1x10<sup>6</sup> HSC-T6 cell (dash lines) and 1 μM Staurosporine (3 hours) treated HSC-T6 cells (full lines) were intracellularly stained with 0.4 μg Cleaved PARP1 Monoclonal Antibody (60555-1-Ig, Clone:4G4C8, red) and Coralite® Plus 647-Goat Anti-Mouse Recombinant Secondary Antibody (H+L)(Cat.NO.RGAM005). Mouse IgG1 isotype control (66360-1-Ig, Clone: 1F8D3, blue) was parallel stained as control. Cells were fixed with 4% PFA. This data was developed using the same antibody




Staurosporine treated and untreated mouse splenocytes were subjected to SDS PAGE followed by western blot with 60555-1-Ig (Cleaved PARP1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with HRP-conjugated Lamin B1 (HRP-66095) antibody as a loading control. This data was developed using the same antibody clone with 60555-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed untreated and 1 μM Staurosporine (3 hours) treated HeLa cells using Cleaved PARP1 antibody (60555-1-Ig, Clone: 4G4C8) at dilution of 1:366 and Multi-rAb Coralite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004). This data was developed using the same antibody clone with 60555-1-PBS in a different storage buffer formulation.



1x10<sup>6</sup> HeLa cell (dash lines) and 1 μM Staurosporine (3 hours) treated HeLa cells (full lines) were intracellularly stained with 0.1 μg Cleaved PARP1 Monoclonal Antibody (60555-1-Ig, Clone:4G4C8, red) and Coralite® Plus 647-Goat Anti-Mouse Recombinant Secondary Antibody (H+L)(Cat.NO.RGAM005). Mouse IgG1 isotype control (66360-1-Ig, Clone: 1F8D3, blue) was parallel stained as control. Cells were fixed with 4% PFA. This data was developed using the same antibody



Staurosporine treated and untreated HSC-T6 cells were subjected to SDS PAGE followed by western blot with 60555-1-Ig (Cleaved PARP1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with HRP-conjugated Lamin B1 (HRP-66095) antibody as a loading control. This data was developed using the same antibody clone with 60555-1-PBS in a different storage buffer formulation.



Cytometric bead array in cell lysate using MP50985-2, Cleaved PARP1 Monoclonal Matched Antibody Pair, PBS Only. Capture antibody: 60555-1-PBS. Detection antibody: 66520-1-PBS. Cell lysate: Non-treated Jurkat and Staurosporine treated Jurkat (10 μg/well). Non-related target Tubulin-beta Recombinant Matched Antibody Pair (MP00550-1) was served as control.