

For Research Use Only

# Anti-Mouse PD-1/CD279 (J43.1)

Catalog Number: 65120-1-Ig



## Basic Information

<b>Catalog Number:</b> 65120-1-Ig	<b>GenBank Accession Number:</b> BC119179	<b>Purification Method:</b> Affinity purification
<b>Size:</b> 500ug, 0.5 mg/ml	<b>GeneID (NCBI):</b> 18566	<b>CloneNo.:</b> J43.1
<b>Source:</b> Armenian Hamster	<b>UNIPROT ID:</b> Q02242	
<b>Isotype:</b> IgG	<b>Full Name:</b> programmed cell death 1	

## Applications

**Tested Applications:**  
FC

**Species Specificity:**  
Mouse

## Background Information

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

## Storage

**Storage:**  
Store at 2-8°C. Stable for one year after shipment.

**Storage Buffer:**  
PBS with 0.09% sodium azide.

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



1x10<sup>6</sup> unstimulated or anti-CD3/CD28 stimulated (2 days) mouse splenocytes were surface stained with 0.5 ug Anti-Mouse PD-1/CD279 (65120-1-Ig, Clone: J43.1) and FITC Goat anti-hamster (Armenian) IgG Antibody at dilution 1:100, and 0.5 ug Coralite® Plus 647 Anti-Mouse CD3 (17A2) (CL647-65077, Clone: 17A2). Cells were not fixed.