

For Research Use Only

# Anti-Human CD49b (AK7)

Catalog Number: 65252-1-Ig



## Basic Information

<b>Catalog Number:</b> 65252-1-Ig	<b>GenBank Accession Number:</b> NM_002203	<b>Purification Method:</b> Affinity purification
<b>Size:</b> 100ug, 0.5 mg/mL	<b>GeneID (NCBI):</b> 3673	<b>CloneNo.:</b> AK7
<b>Source:</b> Mouse	<b>Full Name:</b> integrin, alpha 2 (CD49B, alpha 2 subunit of VLA-2 receptor)	
<b>Isotype:</b> IgG1, kappa	<b>Calculated MW:</b> 126 kDa	

## Applications

**Tested Applications:**  
FC, ELISA

**Species Specificity:**  
Human

## Background Information

CD49b, also known as alpha2 integrin, is a type I transmembrane glycoprotein which forms an adhesion molecule VLA-2 with beta1 integrin. CD49b is a receptor for many (extracellular) matrix and non-matrix molecules (PMID: 23847623). CD49b is expressed on numerous cell types including platelets, activated T cells, B cells, and monocytes. CD49b activation skews NK cell effector function to a phenotype that exhibits elevated IFN- $\gamma$  production with a concomitant inhibition of motility and cytolytic function (PMID: 32971468, 32762681). The Anti-Human CD49b antibody, clone AK7, is derived from Human platelets.

## Storage

**Storage:**  
Store at 2-8°C. Stable for one year after shipment.

**Storage Buffer:**  
PBS with 0.09% sodium azide.

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10<sup>6</sup> human PBMCs were surface stained with 0.2 ug Anti-Human CD49b (65252-1-Ig, Clone:AK7) and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:500. Cells were not fixed. Monocytes were gated.



1X10<sup>6</sup> human PBMCs were surface stained with 0.2 ug Anti-Human CD49b (65252-1-Ig, Clone:AK7) and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:500. Cells were not fixed. Monocytes were gated.