For Research Use Only

Anti-Human IL-1 beta (2A7C6)

Catalog Number: 65276-1-lg



Purification Method:

Affinity purification

CloneNo.:

2A7C6

Basic Information

Catalog Number: 65276-1-Ig

Concentration: 100ug, 1000 µg/ml

Source: Mouse BC008678

GenBank Accession Number:

GeneID (NCBI):

ENSEMBL Gene ID: ENSG00000125538

 Isotype:
 Full Name:

 IgG1
 interleukin 1, beta

 Immunogen Catalog Number:
 Calculated MW:

HZ-1164 269 aa, 31 kDa

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human

Background Information

Interleukin-1 is a pro-inflammatory cytokine with multiple biological effects. The IL-1 gene family encodes three proteins: IL-1 α , IL-1 β and their naturally occurring inhibitor Il-1RN. Interleukin 1 β (IL-1 β), mainly produced by blood monocytes and tissue macrophages, has been implicated in mediating both acute and chronic inflammation. IL-1 β is known to be involved in a variety of cellular activities, including cell proliferation, differentiation and apoptosis. IL-1 β is emerging as a key mediator of carcinogenesis that characterizes host-environment interactions. (PMID: 8630372; 12401481; 23704929; 24618930)


Storage

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1X10^6 LPS and Brefeldin A treated human PBMCs were surface stained with Anti-Human HLA-DR then intracellularly stained with 0.2 ug CoraLite® Plus 488 Anti-Human IL-1 β (CL488-65276, Clone:2A7C6) or Mouse IgG1 Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). Monocytes were gated. This data was developed using the CoraLite ® Plus 488 conjugated antibody of 65276-1-lg.



1X10^6 LPS and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug Coralite® Plus 488 Anti-Human Il-1 β (CL488-65276, Clone:2A7C6) (red) or Mouse IgG1 Isotype Control (blue), or unactivated and Brefeldin A treated human PBMCs were intracellularly stained with 0.2 ug Coralite® Plus 488 Anti-Human Il-1 β (CL488-65276, Clone:2A7C6) (black, dashed line). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).