For Research Use Only

## Anti-Human CD8 (UCHT4) Mouse IgG2a Recombinant Antibody, PBS Only



Catalog Number: 65559-1-PBS

**Basic Information** 

Catalog Number: 65559-1-PBS

Size: 1mg, 2 mg/ml

Source: Mouse Isotype: IgG2a GenBank Accession Number: BC025715

GeneID (NCBI):

925

ENSEMBL Gene ID: ENSG00000153563 UNIPROT ID: P01732

CD8a molecule
Calculated MW:
235 aa, 26 kDa

Full Name:

Purification Method: Protein A purfication

CloneNo.: UCHT4

**Applications** 

**Tested Applications:** 

FC

Species Specificity:

Human

**Background Information** 

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8  $\alpha$  or as a heterodimer of CD8  $\alpha$  and CD8  $\beta$  (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8  $\alpha$   $\beta$  heterodimer while CD8  $\alpha$   $\alpha$  homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR)  $\gamma$   $\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).


Storage

Storage:

Store at -80°C. Stable for one year after shipment.

Storage Buffer: PBS Only

## Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.5 ug Anti-Human CD8 Mouse Recombinant Antibody, PBS Only (65559-1-PBS, Clone:UCHT4) or 0.5 ug Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). Cells were then stained with APC Anti-Human CD3. Cells were not fixed. Lymphocytes were gated.