For Research Use Only

Anti-Human CD54 (15.2) Mouse IgG2a Recombinant Antibody



Purification Method:

Protein A purification

CloneNo.:

15.2

Catalog Number: 65567-1-MR

Basic Information

Catalog Number:

65567-1-MR

Size:

100ug, 1000 $\,\mu$ g/ml

Source: Mouse Isotype:

IgG2a

GenBank Accession Number:

BC015969 GeneID (NCBI):

3383

ENSEMBL Gene ID: ENSG00000090339

Full Name:

intercellular adhesion molecule 1

Calculated MW: 90 kDa

Applications

Tested Applications:

FC

Species Specificity:

Human

Background Information

ICAM-1 (CD54) is a 90-kDa transmembrane glycoprotein of the immunoglobulin superfamily and is critical for the firm attachment and transmigration of leukocytes out of blood vessels and into tissues (PMID: 19307690). ICAM-1 is expressed by several cell types, typically on endothelial cells and cells of the immune system, and its expression can be up-regulated by various stimuli, including TNF- α , INF- γ , IL-1 and thrombin (PMID: 3086451; 9694714; 15979056). It is a ligand for LFA-1 and Mac-1, serves as a receptor for rhinovirus, and is one of several receptors used by Plasmodium falciparum (PMID: 2566624; 2538244; 2475784).

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.125 ug Anti-Human CD54 (15.2) Mouse Recombinant Antibody (65567-1-MR, Clone:15.2) or Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). Cells were then stained with APC Anti-Human CD3 (OKT3) (APC-65133, Clone: OKT3). Cells were not fixed. Lymphocytes were gated.