For Research Use Only

Anti-Rat CD80/B7-1 (3H5) Mouse IgG2a Recombinant Antibody, PBS Only

Catalog Number: 65580-1-PBS



Basic Information

Catalog Number: 65580-1-PBS

GenBank Accession Number: XM_017597885

Purification Method: Protein A purification

Size:

GeneID (NCBI):

CloneNo.:

1mg, 2 mg/ml

25408

3H5

Source: Mouse

Full Name: Cd80 molecule Calculated MW:

Isotype: IgG2a

36 kDa

Applications

Tested Applications:

FC

Species Specificity:

rat

Background Information

CD80 (also known as B7-1) is a type I membrane protein that is a member of the immunoglobulin superfamily, with an extracellular immunoglobulin constant-like domain and a variable-like domain required for receptor binding. It is expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, monocytes, and macrophages. CD80 is the receptor for the proteins CD28 and CTLA-4 found on the surface of T-cells. It is involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation. CD80 also acts as a cellular attachment receptor for adenovirus subgroup B. (PMID: 7545666; 12015893; 16920215)

Storage

Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C


Storage Buffer:

PBS Only

Selected Validation Data



1x10^6 unstimulated or LPS-stimulated rat splenocytes were surface stained with 0.25 ug Anti-Rat CD80/B7-1 Mouse IgG2a Recombinant Antibody (65580-1-MR, Clone: 3H5) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were co-stained with Coralite® Plus 647 Anti-Rat CD45RA (0X-33). Cells were not fixed. This data was developed using the same antibody clone with 65580-1-PBS in a different storage buffer formulation.



65580-1-MR CD80/B7-1(3H5),PE-A

1x10^6 LPS-stimulated rat splenocytes were surface stained with 0.25 ug Anti-Rat CD80/B7-1 Mouse IgG2a Recombinant Antibody (65580-1-MR, Clone: 3H5) (red) or Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4) (blue) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). 1x10^6 unstimulated rat splenocytes were surface stained with 0.25 ug Anti-Rat CD80/B7-1 Mouse IgG2a Recombinant Antibody (65580-1-MR, Clone: 3H5)(black, dashed) and PE-



