For Research Use Only

## Anti-Human CD18 (TS1/18) Mouse IgG2a Recombinant Antibody



**Purification Method:** 

Protein A purification

CloneNo.:

TS1/18

Catalog Number:65595-1-MR

**Basic Information** 

Catalog Number:

65595-1-MR

Size:

100ug, 1000 ug/ml

Source: Mouse Isotype:

IgG2a

GenBank Accession Number: BC005861

GeneID (NCBI):

3689 ENSEMBL Gene ID:

ENSG0000160255 Full Name:

integrin, beta 2 (complement

component 3 receptor 3 and 4 subunit)

Calculated MW: 85 kDa

**Applications** 

**Tested Applications:** 

FC

Species Specificity:

human

## **Background Information**

CD18, also known as integrin beta-2, is a transmembrane protein that forms heterodimers with CD11a, CD11b, CD11c, CD11d (PMID: 1968349). CD11a/CD18 (LFA-1) is expressed by all leukocytes, while CD11b/CD18 (Mac-1) and CD11c/CD18 (p150,95) are normally restricted to expression on monocytes, macrophages, polymorphonuclear lymphocytes (PMNs), and natural killer cells (PMID: 1968349; 3109455; 10946284). CD18 plays an important role in mediating cell adhesion during immune response and defects of CD18 cause leukocyte adhesion deficiency (PMID: 3594570).

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

## **Selected Validation Data**



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD18 Mouse IgG2a Recombinant Antibody (65595-1-MR, Clone: TS1/18) or 0.25 ug Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4) and Coralite® 488-Conjugated AffiniPure Donkey Anti-Mouse IgG(H+L). Cells were then stained with APC Anti-Human CD45. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD18 Mouse IgG2a Recombinant Antibody (65595-1-MR, Clone: TS1/18) (red) or 0.25 ug Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4) (blue) and CoraLite® 488-Conjugated Affini Pure Donkey Anti-Mouse IgG(H+L). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.