Anti-Human CD25 (BC96) Mouse IgG2a Recombinant Antibody


Catalog Number:65620-1-MR



Basic Information	Catalog Number: 65620-1-MR	GenBank Accession Number: BC 156627	Purification Method: Protein A purification
	Size: 100ug, 1 mg/ml	GeneID (NCBI): 3559	CloneNo.: BC96
	Source: Mouse	Full Name: interleukin 2 receptor, alpha	
	Isotype: IgG2a	Calculated MW: 272 aa, 31 kDa	
Applications	Tested Applications: FC		
	Species Specificity: human		
Background Information	Proliferation of T lymphocytes is triggered by the interaction of IL-2 with its specific receptor following T lymphocyte activation. The receptor for IL-2 has three forms, generated by different combinations of three different proteins, the alpha chain (IL-2R alpha), the beta chain (IL-2R beta), and the gamma chain (IL-2R gamma) (PMID: 8476561). IL-2R alpha (also known as CD25) is a type I transmembrane protein present on activated T cells, activated B cells, some thymocytes, myeloid precursors, and oligodendrocytes. CD25 is also found on natural CD4+Foxp3+ Treg cells and a subset of human CD4+ memory T cells (PMID: 22585674).		
Storage	Storage: Store at 2-8°C. Stable for one year after shipment. Storage Buffer: PBS with 0.09% sodium azide.		

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^6 PHA treated human PBMCs were surface stained with 0.25 ug Anti-Human CD25 (BC96) Mouse IgCaa Recombinant Antibody (65620-1-MR, Clone: BC96) (red) or Mouse IgCaa Isotype Control (blue), and PE-conjugated Goat Anti-Mouse IgC. 1x10^6 untreated human PBMCs were surface stained with 0.25 ug Anti-Human CD25 (BC96) Mouse IgCaa Recombinant Antibody (65620-1-MR, Clone: BC96) and PE-conjugated Goat Anti-Mouse IgG (black, dashed). Cells were not

▼