For Research Use Only

Anti-Human CD28 (CD28.2) Mouse IgG2a Recombinant Antibody, PBS Only

Catalog Number:65621-1-PBS




Catalog Number: 65621-1-PBS	GenBank Accession Number: BC093698	Purification Method: Protein A purification
Size: 1mg, 2mg/ml	GenelD (NCBI): 940	CloneNo.: CD28.2
Source: Mouse	ENSEMBL Gene ID: ENSG00000178562	
lsotype: IgG2a	Full Name: CD28 molecule	
	Calculated MW: 220 aa, 25 kDa	
Tested Applications: FC		
Species Specificity: human		
CD28 (T-cell-specific surface glycoprotein CD28), also known as T44 and Tp44, is a 44 kD disulfide-linked homodimeric type I glycoprotein (PMID: 2162180). It is a member of the immunoglobulin superfamily and is expressed on most T lineage cells, NK cell subsets, and plasma cells (PMID: 2162180, 8386518). CD28 may affect in vivo immune responses by functioning both as a cell adhesion molecule linking B and T lymphocytes and as the surface component of a novel signal transduction pathway (PMID: 2162180, 3021470). CD28 binds both CD80 and CD86 with a highly conserved motif MYPPY in the CDR3-like loop (PMID: 15696168, 7964482). CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death (PMID: 1348520).		
Storage: Store at -80°C. The product is shipped with i Storage Buffer: PBS Only	ce packs. Upon receipt, store it immediatel	ly at −80°C
	65621-1-PBS Size: 1mg, 2mg/ml Source: Mouse Isotype: IgG2a Tested Applications: FC Species Specificity: human CD28 (T-cell-specific surface homodimeric type I glycopro expressed on most T lineage vivo immune responses by fi surface component of a nove CD86 with a highly conserve major co-stimulatory molecu (PMID: 1348520). Storage: Stora at -80°C. The product is shipped with i Storage Buffer:	65621-1-PBS BC093698 Size: GeneID (NCBI): 1mg, 2mg/ml 940 Source: ENSEMBL Gene ID: Mouse ENSG0000178562 Isotype: Full Name: IgG2a CD28 molecule Calculated MW: 220 aa, 25 kDa Tested Applications: FC Species Specificity: human CD28 (T-cell-specific surface glycoprotein CD28), also known as T44 an homodimeric type I glycoprotein (PMID: 2162180). It is a member of the expressed on most T lineage cells, NK cell subsets, and plasma cells (PV vivo immune responses by functioning both as a cell adhesion molecul surface component of a novel signal transduction pathway (PMID: 2162 CD86 with a highly conserved motif MYPPY in the CDR3-like loop (PMID major co-stimulatory molecule, inducing T lymphocyte activation and I (PMID: 1348520). Storage: Storage Store at -80°C. The product is shipped with ice packs. Upon receipt, store it immediated Storage Buffer:


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^{^6} human PBMCs were surface stained with 0.25 ug Anti-Human CD28 (CD28.2) Mouse IgG2a Recombinant Antibody (65621-1-MR, Clone: CD28.2) or Mouse IgG2a Isotype Control (C1.18.4), (65208-1-Ig, Clone: C1.18.4), and PE-conjugated Goat Anti-Mouse IgG. Cells were co-stained with APC Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody. Cells were not fixed. This data was developed using the same antibody clone with 65621-1-PBS in a different storage buffer formulation. ▼



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD28 (CD28.2) Mouse IgG2a Recombinant Antibody (65621-1-MR, Clone: CD28.2) (red) or Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) (blue), and PE-conjugated Goat Anti-Mouse IgG. Cells were not fixed. This data was developed using the same antibody clone with 65621-1-PBS in a different storage buffer formulation.