For Research Use Only

Anti-Human CD8 (SK1) Mouse IgG2a Recombinant Antibody, PBS Only

Catalog Number:65630-1-PBS




Basic Information	Catalog Number: 65630-1-PBS	GenBank Accession Number: BC025715	Purification Method: Protein A purification
	Size: 1mg, 2mg/ml	Genel D (NCBI): 925	CloneNo.: SK1
	Source: Mouse	ENSEMBL Gene ID: ENSG00000153563	
	Isotype: IgG2a	Full Name: CD8a molecule	
		Calculated MW: 235 aa, 26 kDa	
Applications	Tested Applications: FC		
	Species Specificity: human		
Background Information	CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).		
Storage	Storage: Store at -80°C. The product is shipped with ice par Storage Buffer: PBS Only	cks. Upon receipt, store it immediatel	yat-80°C


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^{^6} human PBMCs were surface stained with 0.25 ug Anti-Human CD8 (SK1) Mouse IgG2a Recombinant Antibody (65630-1-MR, Clone: SK1) or 0.25 ug Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4), and PEconjugated Goat Anti-Mouse IgG. Cells were then stained with APC Anti-Human CD3. Cells were not fixed. Lymphocytes were gated. This data was developed using the same antibody clone with 65630-1-PBS in a different storage buffer formulation.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD8 (SK1) Mouse IgG2a Recombinant Antibody (65630-1-MR, Clone: SK1) (red) or 0.25 ug Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4) (blue), and PEconjugated Goat Anti-Mouse IgG. Cells were not fixed. Lymphocytes were gated. This data was developed using the same antibody clone with 65630-1-PBS in a different storage buffer formulation.