

For Research Use Only

P4HA1 Monoclonal antibody, PBS Only



Catalog Number: 66101-1-PBS

Featured Product

Basic Information

Catalog Number:

66101-1-PBS

Size:

1mg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG17867

GenBank Accession Number:

BC034998

GeneID (NCBI):

5033

UNIPROT ID:

P13674

Full Name:

prolyl 4-hydroxylase, alpha polypeptide I

Calculated MW:

534 aa, 61 kDa

Observed MW:

61 kDa

Purification Method:

Protein A purification

CloneNo.:

1H10A6

Applications

Tested Applications:

WB, IHC, Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

P4HA1 (Prolyl 4-hydroxylase subunit alpha-1) is also named as P4HA and belongs to the P4HA family, which play a central role in collagen synthesis. P4HA1 catalyzes the post-translational formation of 4-hydroxyproline in -Xaa-Pro-Gly- sequences in collagens and other proteins. The gene encodes a polypeptide of 517 amino acid residues and a signal peptide of 17 amino acids and the full length protein has two glycosylation sites (uniprot). It can exist as a heteromer, dimer or tetramer (refer to: GENATLAS). P4HA1 has 3 isoforms produced by alternative splicing with the molecular weight of 61 kDa, 61 kDa and 59 kDa. This antibody is specific to P4HA1.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66101-1-Ig (P4HA1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66101-1-PBS in a different storage buffer formulation.



PC-3 cells were subjected to SDS PAGE followed by western blot with 66101-1-Ig (P4HA1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66101-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66101-1-Ig (P4HA1 Antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66101-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66101-1-Ig (P4HA1 Antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66101-1-PBS in a different storage buffer formulation.