

For Research Use Only

CSK Monoclonal antibody

Catalog Number: 66115-1-Ig **1 Publications**



Basic Information

Catalog Number: 66115-1-Ig	GenBank Accession Number: BC106073	Purification Method: Protein A purification
Size: 1900 µg/ml	GeneID (NCBI): 1445	CloneNo.: 3B12C5
Source: Mouse	UNIPROT ID: P41240	Recommended Dilutions: WB 1:500-1:2000 IF/ICC 1:20-1:200
Isotype: IgG2b	Full Name: c-src tyrosine kinase	
Immunogen Catalog Number: AG12103	Calculated MW: 450 aa, 50 kDa	
	Observed MW: 50 kDa	

Applications

Tested Applications: IF/ICC, WB, ELISA	Positive Controls: WB : Jurkat cells, HeLa cells, NIH/3T3 cells, RAW 264.7 cells IF/ICC : A431 cells,
Cited Applications: WB	
Species Specificity: human, mouse	
Cited Species: mouse	

Background Information

CSK(Tyrosine-protein kinase CSK) is also named as C-Src kinase, protein-tyrosine kinase CYL and belongs to the protein kinase superfamily. Csk is a non-receptor tyrosine kinase with a molecular mass of 50 kDa. It contains the SH3 and SH2 domains in its N-terminus and a kinase domain in its C-terminus. It negatively regulates the kinase activity of SFKs and plays an important role in physiological functions via signaling pathways for cell proliferation, differentiation, adhesion, and migration(PMID:23548896). Csk may play a role in tumor suppression by inhibiting the oncogenic activity of SFKs. Indeed, overexpression of Csk can inhibit tumor growth of human colon cancer cells(PMID:23139636). This protein can be autophosphorylated(PMID:11181701).

Notable Publications

Author	Pubmed ID	Journal	Application
Jiandong Sun	34185441	Andrology	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 66115-1-Ig (CSK antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of A431 cells using 66115-1-Ig (CSK antibody) at dilution of 1:50 and Rhodamine-Goat anti-Mouse IgG.