For Research Use Only

ERK1/2 Monoclonal antibody

Catalog Number: 66192-1-lg 26 Publications



Basic Information

Catalog Number: 66192-1-lg Size:

1240 µg/ml
Source:
Mouse
Isotype:
IgG1

Immunogen Catalog Number:

AG20350

G20350

GenBank Accession Number:

BC013992 GeneID (NCBI): 5595 UNIPROT ID: P27361

Full Name: mitogen-activated protein kinase 3

43 kDa Observed MW: 38-44 kDa

Calculated MW:

Positive Controls:

WB: HeLa cells, HepG2 cells, MCF-7 cells, HEK-293 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells

Purification Method:

Protein G purification

Recommended Dilutions:

WB 1:2000-1:10000 IHC 1:2000-1:6000

IF 1:200-1:800

CloneNo.:

4G8A11

 $\hbox{IHC:} \hbox{ human ovary tumor tissue, human cervical cancer}\\$

tissue

IF: HeLa cells,

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications: CoIP, WB Species Specificity:

human, mouse
Cited Species:
human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

ERK1 and ERK2 belongs to the protein kinase superfamily. It is involved in both the initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors such as ELK-1. ERK1/2 catalized the reaction: ATP + a protein = ADP + a phosphoprotein. It is activated by tyrosine phosphorylation in response to INS and NGF. This antibody can recognize both ERK1 and ERK2 with the molecular mass of 38-44 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Li Li	36166754	Br J Pharmacol	WB
Zhichao Dou	32956704	Exp Cell Res	WB
Mengnan Hu	36064579	Oncogene	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66192-1-1g (ERK1/2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 66192-1-Ig (ERK1/2 antibody) at dilution of 1:4000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 66192-1-Ig (ERK1/2 antibody) at dilution of 1:4000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using ERK1/2 antibody (66192-1-lg, Clone: 4G8A11) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse lgG(H+L).