

# FAK Monoclonal antibody

Catalog Number: 66258-1-Ig

Featured Product

35 Publications

## Basic Information

## Catalog Number:

66258-1-Ig

## Size:

2592 µg/ml

## Source:

Mouse

## Isotype:

IgG2c

## Immunogen Catalog Number:

AG17966

## GenBank Accession Number:

BC028733

## GeneID (NCBI):

5747

## UNIPROT ID:

Q05397

## Full Name:

PTK2 protein tyrosine kinase 2

## Calculated MW:

1052 aa, 119 kDa

## Observed MW:

110 kDa

## Purification Method:

Protein A purification

## CloneNo.:

2C5B9

## Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:50-1:500

IF 1:200-1:800

## Applications

## Tested Applications:

IF/ICC, IHC, WB, ELISA

## Cited Applications:

IF, IP, WB

## Species Specificity:

human, mouse, rat

## Cited Species:

human, rat, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

**WB** : hTERT-RPE1 cells, human testis tissue, MCF-7 cells, HCT 116 cells, NIH/3T3 cells, RAW 264.7 cells, ROS1728 cells, HepG2 cells, HeLa cells, Jurkat cells, K-562 cells, pig brain tissue, rabbit brain tissue, rat brain tissue, mouse brain tissue

**IHC** : human breast cancer tissue,

**IF** : HUVEC cells, A549 cells

## Background Information

FAK (Focal adhesion kinase 1) is also named as FAK1, FADK, pp125FAK, FAK and belongs to the protein kinase superfamily. It is a critical tyrosine kinase that modulates cell adhesion, migration, proliferation and survival in response to extracellular signals (PMID:19664602). It also acts as a pivotal signal 'integrator', controlling and coordinating cellular responses that include cell migration, survival, proliferation and, epithelial tissue repair after DNA damage (PMID:20966971). This protein has some isoforms produced by alternative promoter usage and alternative splicing, and the range of the molecular weights are 100-120kD and 40-60kD.

## Notable Publications

Author	Pubmed ID	Journal	Application
Jiao Wu	34517345	Aging (Albany NY)	WB
Olga Alekhina	27789711	J Biol Chem	WB
Andreas Ritter	31640218	Cells	IF

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




WB result of FAK antibody (66258-1-Ig; 1:6000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FAK transfected MCF-7 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 66258-1-Ig (FAK antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66258-1-Ig (FAK antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66258-1-Ig (FAK antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HUVEC cells using FAK antibody (66258-1-Ig, Clone: 2C5B9) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).