## For Research Use Only

## OGDH Monoclonal antibody, PBS Only



Catalog Number: 66285-1-PBS

Basic Information

Catalog Number:

Size: 1 mg/ml

66285-1-PBS

BC004964 Genel D (NCBI): 4967

Source: UNIPROT ID:
Mouse Q02218

Isotype: Full Name:
IgG2b oxoglutarate (alpha-ketoglutarate)

Immunogen Catalog Number: dehydrogenase (lipoamide)

AG7484

Calculated MW: 116 kDa

GenBank Accession Number:

Observed MW: 116 kDa

**Applications** 

Tested Applications: WB,Indirect ELISA,IHC,IF Species Specificity: human, rat

**Background Information** 


OGDH(2-oxoglutarate dehydrogenase, mitochondrial) is also named as OGDC-E1and belongs to the alphaketoglutarate dehydrogenase family. It is one of the three enzymatic components of 2-oxoglutarate dehydrogenase complex,which catalyzes the overall conversion of 2-oxoglutarate to succinyl-CoA and CO2. This protein has 3 isoforms produced by alternative splicing.

Storage

Storage: Store at -80°C. Storage Buffer: PBS Only Purification Method: Protein A purification

CloneNo.: 1H7A2


## Selected Validation Data




human heart tissue were subjected to SDS PAGE followed by western blot with 66285-1-lg (OGDH Antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66285-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66285-1-lg (OGDH Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66285-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66285-1-lg (OGDH Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66285-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using OGDH antibody (66285-1-Ig, Clone: 1H7A2) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-phalloidin (red). This data was developed using the same antibody clone with 66285-1-PBS in a different storage buffer formulation.