For Research Use Only

## IL-12/IL-23 p40 Monoclonal antibody, PBS Only



**Purification Method:** 

Protein A purification

CloneNo.:

4B2A8

Catalog Number: 66288-1-PBS

**Basic Information** 

Catalog Number: 66288-1-PBS

Size:
1 mg/ml
Source:
Mouse
Isotype:

Immunogen Catalog Number:

AG21429

IgG2a

GenBank Accession Number:

BC067498 GeneID (NCBI): 3593 UNIPROT ID:

P29460

Full Name:

interleukin 12B (natural killer cell stimulatory factor 2, cytotoxic lymphocyte maturation factor 2, p40)

Calculated MW: 328 aa, 37 kDa Observed MW: 37 kDa

**Applications** 

Tested Applications: WB,Indirect ELISA Species Specificity: human, pig

## **Background Information**

IL-12 and IL-23 are heterodimeric cytokines that share a common p40 subunit (PMID: 11114383). IL-12 is composed of the IL-12 p40 subunit linked to the IL-12 p35 subunit, and the heterodimer signals through the IL-12 receptor (IL-12R), which comprises the IL-12R  $\beta$  1 and IL-12R  $\beta$  2 subunits. IL-23 is composed of the IL-23 p19 subunit and the IL-12 p40 (IL-12/23p40) subunit, which signals through IL-23R and IL-12R  $\beta$  1 (PMID: 11114383; 26121196). IL-12/IL-23 p40 also exists as a monomer and as a homodimer which can act as a potent IL-12 antagonist (PMID: 8958912; 18783467). IL-12/IL-23 p40 is produced by antigen-presenting cells, such as dendritic cells (DCs), monocytes, macrophages, neutrophils and, to a lesser extent, B cells (PMID: 20476918).

Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

## Selected Validation Data



human liver tissue were subjected to SDS PAGE followed by western blot with 66288-1-lg (IL12B Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66288-1-PBS in a different storage buffer formulation.