

For Research Use Only

# CD27 Monoclonal antibody, PBS Only

Catalog Number: 66308-1-PBS



## Basic Information

<b>Catalog Number:</b> 66308-1-PBS	<b>GenBank Accession Number:</b> BC012160	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 1mg/ml	<b>GeneID (NCBI):</b> 939	<b>CloneNo.:</b> 1C1G3
<b>Source:</b> Mouse	<b>ENSEMBL Gene ID:</b> ENSG00000139193	
<b>Isotype:</b> IgG1	<b>UNIPROT ID:</b> P26842	
<b>Immunogen Catalog Number:</b> AG24537	<b>Full Name:</b> CD27 molecule	
	<b>Calculated MW:</b> 29 kDa	
	<b>Observed MW:</b> 50-55 kDa	

## Applications

**Tested Applications:**  
WB, IHC, IF-P, FC, ELISA

**Species Specificity:**  
human, pig

## Background Information

CD27 (also known as TNFRSF7) is a type I glycoprotein expressed on some B cells and the majority of T cells, and is a member of the tumor necrosis factor (TNF) receptor family. CD27 is required for generation and long-term maintenance of T cell immunity (PMID: 11062504). It is a receptor for CD70 (CD27L). Ligation of CD27 by CD70 induces strong ubiquitination of TRAF and the activation of both canonical and non-canonical NF-kappaB pathways, as well as the JNK pathway (PMID: 19426224). CD27 may also play a role in apoptosis through association with SIVA1.

## Storage

**Storage:**  
Store at -80°C.  
**The product is shipped with ice packs. Upon receipt, store it immediately at -80°C**

**Storage Buffer:**  
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66308-1-Ig (CD27 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66308-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66308-1-Ig (CD27 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66308-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66308-1-Ig (CD27 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66308-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using 66308-1-Ig (CD27 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66308-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using 66308-1-Ig (CD27 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66308-1-PBS in a different storage buffer formulation.



$1 \times 10^6$  human PBMCs were surface stained with  $0.2 \mu\text{g}$  CD27 Monoclonal antibody (66308-1-Ig, Clone:1C1G3) or  $0.2 \mu\text{g}$  Mouse IgG1 isotype control (66360-1-Ig, Clone: T1F8D3F10), and CoraLite® Plus 647-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO.RGAM005). Cells were not fixed. This data was developed using the same antibody clone with 66308-1-PBS in a different storage buffer formulation.