For Research Use Only

Gamma Tubulin Monoclonal antibody, PBS Only

Antibodies | ELISA kits | Proteins WWW.ptglab.com

Catalog Number:66320-1-PBS

Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 66320-1-PBS BC000619 Protein A purification GeneID (NCBI): CloneNo.: Size: 1 mg/ml 7283 3F9H8 Source: UNIPROT ID: Mouse P23258 Full Name: Isotype: lgG2a tubulin, gamma 1 Calculated MW: Immunogen Catalog Number: AG23913 51 kDa **Observed MW:** 48-55 kDa **Applications Tested Applications:** WB, IF, IHC, Indirect ELISA Species Specificity: human, Canine, mouse, rat **Background Information** Gamma tubulin is a member of tubulin superfamily and is a key component required for microtubule nucleation and

Gamma tubulin is a member of tubulin superfamily and is a key component required for microtubule nucleation and stabilization. It is concentrated at the pericentriolar material of centrosomes in interphase cells, predominantly on spindle poles in mitotic cells, while found in midbodies during cytokinesis. Overexpression of gamma tubulin has been found in various cancers including breast cancer and gliomas. This antibody can well label the centrosome structure.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunofluorescent analysis of (-20°C Ethanol) fixed MDCK cells using 66320-1-Ig (gamma tubulin antibody) at dilution of 1:200 and Coralite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66320-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed MDCK cells using 66320-1-lg (gamma tubulin antibody) at dilution of 1:200 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse lgG(H+L). This data was developed using the same antibody clone with 66320-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using Gamma Tubulin antibody (66320-1-lg, Clone: 3F9H8) at dilution of 1:1000 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), Alpha Tubulin antibody (11224-1-AP, red). This data was developed using the same antibody clone with 66320-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66320-1-Ig (gamma tubulin antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66320-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66320-1-1g (gamma tubulin antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66320-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66320-1-Ig (gamma tubulin antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66320-1-PBS in a different storage buffer formulation.