For Research Use Only

## D2HGDH Monoclonal antibody, PBS Only



Catalog Number: 66364-1-PBS

**Basic Information** 

Catalog Number: 66364-1-PBS

BC036604

Protein G purification

Size: 1mg/ml GeneID (NCBI): 728294

GenBank Accession Number:

CloneNo.: 1E9B3

**Purification Method:** 

Source: **UNIPROT ID:** Mouse Q8N465 Full Name: Isotype:

lgG1 D-2-hydroxyglutarate dehydrogenase

Calculated MW: Immunogen Catalog Number:

AG5036 521 aa, 56 kDa

Observed MW: 56 kDa

**Applications** 

**Tested Applications:** 

Indirect ELISA, IF/ICC, IHC, WB

Species Specificity: rat, mouse, human

**Background Information** 

 $\label{lem:decomposition} D2 HGDH (D-2-hydroxyglutarate\ dehydrogenase,\ mit ochondrial)\ is\ also\ named\ as\ D2 HGD\ and\ belongs\ to\ the\ FAD-particles and\ belongs\ to\ the\ particles are also\ named\ as\ D2 HGD\ and\ belongs\ to\ the\ particles are also\ named\ as\ D2 HGD\ and\ belongs\ to\ the\ particles are also\ named\ as\ D2 HGD\ and\ belongs\ to\ the\ particles are also\ named\ as\ D2 HGD\ and\ belongs\ to\ the\ particles are also\ named\ as\ D2 HGD\ and\ belongs\ to\ the\ particles are also\ named\ as\ D2 HGD\ and\ belongs\ to\ the\ particles are also\ named\ as\ D2 HGD\ and\ belongs\ to\ the\ particles are also\ named\ as\ D2 HGD\ and\ belongs\ to\ the\ particles are also\ named\ named$ binding oxidoreductase/transferase type 4 family.It catalyzes the oxidation of D-2-hydroxyglutarate to alphaketoglutarate. Defects in D2HGDH are the cause of D-2-hydroxyglutaric aciduria type 1 (D2HGA1).

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS Only


## Selected Validation Data




1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human D2HGDH (66364-1-1g, Clone:1E9B3) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66364-1-PBS in a different storage buffer formulation.




NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 66364-1-lg (D2HGDH Antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66364-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66364-1-lg (D2HGDH Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66364-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66364-1-lg (D2HGDH Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66364-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using D2HGDH antibody (66364-1-lg, Clone: 1E9B3) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red). This data was developed using the same antibody clone with 66364-1-PBS in a different storage buffer formulation.