For Research Use Only

Thioredoxin Monoclonal antibody, PBS Only



Catalog Number:66475-1-PBS

Featured Product

Basic Information

Catalog Number:

GenBank Accession Number: BC003377

Purification Method: Protein A purification

66475-1-PBS

GeneID (NCBI):

CloneNo.:

Size: 1 mg/ml

7295

1F3C1

Source:

UNIPROT ID: P10599

Mouse Isotype: lgG2a

Full Name: thioredoxin

Immunogen Catalog Number: AG6355

Calculated MW: 12 kDa

12 kDa

Observed MW:

Applications

Tested Applications:

WB,Indirect ELISA,IHC,IF

Species Specificity:

Human

Background Information


TXN, TRDX, TRX1, ADF and SASP, belongs to the thioredoxin family. It participates in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyzes dithiol-disulfide exchange reactions. TXN plays a role in the reversible S-nitrosylation of cysteine residues in target proteins, and thereby contributes to the response to intracellular nitric oxide. Nitrosylates the active site Cys of CASP3 in response to $nitric\ oxide\ (NO), and\ thereby\ inhibits\ caspase-3\ activity.\ TXN\ induces\ the\ FOS/JUN\ AP-1\ DNA-binding\ activity\ in$ ionizing radiation (IR) cells through its oxidation/reduction status and stimulates AP-1 transcriptional activity.

Storage

Store at -80°C.

Storage Buffer: PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66475-1-1g (TXN antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 66475-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 66475-1-Ig (Thioredoxin antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66475-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Thioredoxin antibody (66475-1-Ig, Clone: 1F3C1) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66475-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66475-1-1g (Thioredoxin antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66475-1-PBS in a different storage buffer formulation



WB result of Thioredoxin antibody (66475-1-1g; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Thioredoxin transfected HeLa cells. This data was developed using the same antibody clone with 66475-1-PBS in a different storage buffer formulation.