For Research Use Only

SMARCA4/BRG1 Monoclonal antibody, PBS Only



Purification Method:

Protein A purification

CloneNo.:

2E6B6

Catalog Number:66561-1-PBS

Featured Product

Basic Information

Catalog Number: 66561-1-PBS

Size:
1 mg/ml
Source:
Mouse
Isotype:

Immunogen Catalog Number:

AG16256

lgG2b

GenBank Accession Number:

BC150298 GeneID (NCBI): 6597

UNIPROT ID:

P51532

Full Name: SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 4

Calculated MW: 1647 aa, 185 kDa Observed MW: 185 kDa

Applications

Tested Applications: WB,IP,IF,ELISA Species Specificity: Human, rat


Background Information

SMARCA4, also named as BAF190A, BRG1, SNF2B and SNF2L4, belongs to the SNF2/RAD54 helicase family. SMARCA4 is a transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. It is a component of the CREST-BRG1 complex, a multiprotein complex that regulates promoter activation by orchestrating a calcium-dependent release of a repressor complex and a recruitment of an activator complex. It is also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene.


Storage

Storage: Store at -80°C. Storage Buffer: PBS only


Selected Validation Data




IP result of anti-SMARCA4/BRG1 (IP:66561-1-lg, 5ug; Detection:66561-1-lg 1:3000) with HeLa cells lysate 3200 ug. This data was developed using the same antibody clone with 66561-1-PBS in a different storage buffer formulation.




WB result of SMARCA4/BRG1 antibody (66561-1-lg; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SMARCA4/BRG1 transfected HepG2 cells. This data was developed using the same antibody clone with 66561-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 66561-1-1g (SMARCA4/BRG1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66561-1-PBS in a different storage buffer formulation.




COLO 320 cells were subjected to SDS PAGE followed by western blot with 66561-1-lg (SMARCA4/BRG1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66561-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SMARCA4/BRG1 antibody (66561-1-lg, Clone: 2E6B6) at dilution of 1:50000 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 66561-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed PC-3 cells using SMARCA4/BRG1 antibody (66561-1-lg, Clone: 2E6B6) at dilution of 1:50000 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 66561-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SMARCA4/BRG1 antibody (66561-1-1g, Clone: 2E6B6) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 66561-1-PBS in a different storage buffer formulation.