

For Research Use Only

# CD100 Monoclonal antibody, PBS Only



Catalog Number: 66582-1-PBS

Featured Product

## Basic Information

Catalog Number:

66582-1-PBS

Size:

1 mg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG26461

GenBank Accession Number:

BC054500

GeneID (NCBI):

10507

UNIPROT ID:

Q92854

Full Name:

sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D

Calculated MW:

96 kDa

Observed MW:

150 kDa

Purification Method:

Protein A purification

CloneNo.:

1C1A6

## Applications

Tested Applications:

WB, Indirect ELISA, IHC

Species Specificity:

Human

## Background Information

CD100 (also known as SEMA4D), a 150 kDa homodimer, belongs to the family of immune semaphorins, and CD100 is abundantly expressed on resting T cells, NK cells, and antigen-presenting cells. As a transmembrane glycoprotein, it is digested to its soluble form, sCD100, by specific matrix metalloproteinases. CD100 has been reported that the expression levels of CD100, and one of its receptors, plexin-B1 (PLXNB1), are increased in head and neck, prostate, colon, breast, and lung cancers, and the interaction between them provides oncogenic signaling essential for tumor growth and metastasis.

## Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




various cells were subjected to SDS PAGE followed by western blot with 66582-1-Ig (CD100 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66582-1-PBS in a different storage buffer formulation.



WB result of CD100 antibody (66582-1-Ig; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-CD100 transfected HeLa cells. This data was developed using the same antibody clone with 66582-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66582-1-Ig (CD100 antibody) at dilution of 1:450 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66582-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66582-1-Ig (CD100 antibody) at dilution of 1:450 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66582-1-PBS in a different storage buffer formulation.