

For Research Use Only

# c-Fos Monoclonal antibody, PBS Only

Catalog Number: 66590-1-PBS

Featured Product

1 Publications



## Basic Information

Catalog Number:

66590-1-PBS

Size:

1mg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG24340

GenBank Accession Number:

BC004490

GeneID (NCBI):

2353

UNIPROT ID:

P01100

Full Name:

FOS

Calculated MW:

41 kDa

Observed MW:

55-60 kDa

Purification Method:

Protein G purification

CloneNo.:

1G2C5

## Applications

Tested Applications:

WB, Indirect ELISA

Cited Applications:

IHC

Species Specificity:

human, mouse, rat

Cited Species:

human

## Background Information

c-Fos, also named as FOS and G0/G1 switch regulatory protein 7, is a 380 amino acid protein, which contains 1 bZIP (basic-leucine zipper) domain and belongs to the bZIP family. c-Fos is expressed at very low levels in quiescent cells. When cells are stimulated to reenter growth, c-Fos undergo 2 waves of expression, the first one peaks 7.5 minutes following FBS induction. At this stage, the c-Fos protein is localized endoplasmic reticulum. The second wave of expression occurs at about 20 minutes after induction and peaks at 1 hour. At this stage, the c-FOS protein becomes nuclear. c-Fos is a very short-lived intracellular protein, which is very easy to degrade. The calculated molecular weight of c-Fos is 40 kDa, but Phosphorylated c-Fos protein is about 60-65 kDa. It is involved in important cellular events, including cell proliferation, differentiation and survival; genes associated with hypoxia; and angiogenesis; which makes its dysregulation an important factor for cancer development. It can also induce a loss of cell polarity and epithelial-mesenchymal transition, leading to invasive and metastatic growth in mammary epithelial cells. Expression of c-Fos is an indirect marker of neuronal activity because c-Fos is often expressed when neurons fire action potentials. Upregulation of c-Fos mRNA in a neuron indicates recent activity.

## Notable Publications

Author	Pubmed ID	Journal	Application
Hongtao Zhu	39447031	Neuro Oncol	IHC

## Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



WB result of c-Fos antibody (66590-1-Ig; 1:40000; incubated at room temperature for 1.5 hours) with sh-Control and sh-c-Fos transfected HepG2 cells. This data was developed using the same antibody clone with 66590-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66590-1-Ig (c-Fos antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66590-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66590-1-Ig (c-Fos antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66590-1-PBS in a different storage buffer formulation.