

For Research Use Only

# GORASP2 Monoclonal antibody, PBS Only



Catalog Number: 66627-1-PBS

## Basic Information

|   |  |   |
|---|--|---|
| <b>Catalog Number:</b><br>66627-1-PBS       | <b>GenBank Accession Number:</b><br>BC007770                       | <b>Purification Method:</b><br>Protein G purification |
| <b>Size:</b><br>1 mg/ml                     | <b>GeneID (NCBI):</b><br>26003                                     | <b>CloneNo.:</b><br>1C9A3                             |
| <b>Source:</b><br>Mouse                     | <b>UNIPROT ID:</b><br>Q9H8Y8                                       |   |
| <b>Isotype:</b><br>IgG1                     | <b>Full Name:</b><br>golgi reassembly stacking protein 2,<br>55kDa |   |
| <b>Immunogen Catalog Number:</b><br>AG17607 | <b>Calculated MW:</b><br>55 kDa                                    |   |
|   | <b>Observed MW:</b><br>59 kDa                                      |   |

## Applications

**Tested Applications:**  
WB, Indirect ELISA, IHC, IF

**Species Specificity:**  
Human, Rat, Pig

## Background Information

GORASP2 (Golgi reassembly-stacking protein 2, or GRS2) has multiple phosphorylation sites and can be phosphorylated by MKK/ERK pathway during mitosis. GORASP2 plays a role in the assembly and membrane stacking of the Golgi cisternae, and in the process by which Golgi stacks reform after mitotic breakdown. GORASP2 is supposed to regulate the intracellular transport and presentation of a defined set of transmembrane proteins, such as transmembrane TGFA.

## Storage

**Storage:**  
Store at -80°C.

**Storage Buffer:**  
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)




W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66627-1-Ig (GORASP2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66627-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using GORASP2 antibody (66627-1-Ig, Clone: 1C9A3) at dilution of 1:1000 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66627-1-PBS in a different storage buffer formulation.