For Research Use Only

WNT2 Monoclonal antibody

Catalog Number:66656-1-lg Featured Product

5 Publications



Basic Information

Catalog Number: 66656-1-lg Size:

1000 µg/ml Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG25185

Applications Tested Applications: IF/ICC, WB, ELISA

Cited Applications:

Species Specificity: Human, mouse, rat **Cited Species:** human, rat

GenBank Accession Number:

BC029854 GeneID (NCBI): 7472 **UNIPROT ID:**

P09544 Full Name:

wingless-type MMTV integration site family member 2

Calculated MW: 40 kDa Observed MW: 36 kDa

Positive Controls:

WB: HeLa cells, HSC-T6 cells, Jurkat cells, RAW 264.7 cells, rat brain tissue, NIH/3T3 cells, mouse brain tissue, HEK-293 cells, MCF-7 cells

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000 IF 1:50-1:500

CloneNo.:

3H7B10

IF: MCF-7 cells,

Background Information

Wnt2 is a secreted glycoprotein that is one of the canonical Wnt ligands. The WNT gene family consists of $structurally\ related\ genes\ which\ encode\ secreted\ signaling\ proteins.\ These\ proteins\ have\ been\ implicated\ in$ oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. Wnt2 is known to directly regulate β -catenin activity, the key player in proliferation and inflammation. Recently, studies have demonstrated that Wnt2 is overexpressed in various human cancers, including esophageal, gastric, colorectal, breast, lung, cervical, and malignant glioma.

Notable Publications

Author	Pubmed ID	Journal	Application
Meizhong Peng	36237833	Int J Endocrinol	WB
Zewu Zhu	34724098	Cell Mol Life Sci	IF
Yuan Qu	30463708	J Surg Res	WB


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66656-1-lg (WNT2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using 66656-1-1g (WNT2 antibody) at dilution of 1:100 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).