For Research Use Only

OCIAD1 Monoclonal antibody, PBS Only (Capture)

www.ptglab.com

Catalog Number: 66698-1-PBS

Featured Product

Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purification

66698-1-PBS

GeneID (NCBI):

UNIPROT ID:

Q9NX40

BC003409

CloneNo.:

Size: 1mg/ml

IgG2a

AG9977

54940

1C10C3

Source: Mouse Isotype:

OCIA domain containing 1

Immunogen Catalog Number:

Calculated MW:

Full Name:

28 kDa

Observed MW:

35 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, Cytometric bead array, Indirect

Species Specificity:

human, mouse, rat

Background Information

OCIAD1 was first identified by immunoscreening of an ovarian carcinoma cDNA expression library with ascites fluid from ovarian cancer patients (PMID: 11162530). OCIAD1 has been reported as a key player in ovarian cancer cell adhesion, as well as a key player in generating ovarian cancer recurrence (PMID: 18328549; 20515946). In addition to its roles in cancer, OCIAD1 participates in maintaining stem cell potency by regulating the Jak/STAT pathway (PMID: 23972987). Several alternatively spliced forms of OCIAD1 gene have been identified. The longest form (1.4 kb) is predicted to encode for a 27.6 kDa protein of 245 amino acids. This antibody detects OCIAD1 with an apparent molecular weight of ~35 kDa as has been demonstrated by several researches (PMID: 27345969; 27345976).

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

PBS Only


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 66698-1-Ig (OCIAD1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66698-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66698-1-lg (OCIAD1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66698-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66698-1-lg (OCIAD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66698-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using OCIAD1 antibody (66698-1-Ig, Clone: 1C10C3) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66698-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using OCIAD1 antibody (66698-1-Ig, Clone: 1C10C3) at dilution of 1:400 and Coralite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66698-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using OCIAD1 antibody (66698-1-1g, Clone: 1C10C3) at dilution of 1:400 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red). This data was developed using the same antibody clone with 66698-1-PBS in a different storage buffer formulation.



Cytometric bead array standard curve of MP50614-1, OCIAD1 Monoclonal Matched Antibody Pair, PBS Only. Capture antibody: 66698-1-PBS. Detection antibody: 66698-2-PBS. Standard:Ag9977. Range: 0.098-100 ng/mL



Cytometric bead array standard curve of MP50614-2, OCIAD1 Monoclonal Matched Antibody Pair, PBS Only. Capture antibody: 66698-1-PBS. Detection antibody: 66698-3-PBS. Standard:Ag9977. Range: 0.098-100 ng/mL