

For Research Use Only

TRAF4 Monoclonal antibody, PBS Only



Catalog Number: 66755-1-PBS

Featured Product

Basic Information

Catalog Number:

66755-1-PBS

Size:

1 mg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG27221

GenBank Accession Number:

BC001769

GeneID (NCBI):

9618

UNIPROT ID:

Q9BUZ4

Full Name:

TNF receptor-associated factor 4

Calculated MW:

54 kDa

Observed MW:

50-54 kDa

Purification Method:

Protein A purification

CloneNo.:

2C9A11

Applications

Tested Applications:

WB, Indirect ELISA, IHC, IF

Species Specificity:

Human, pig

Background Information

The tumor necrosis factor receptor-associated factor (TRAF) family, consisting of seven members (TRAF1-7), is a group of signaling adaptors which participate in various signaling pathways through binding of the tumor necrosis factor receptor superfamily. As one of the important member of TRAF family proteins, TRAF4 was initially isolated from breast carcinomas and identified as the first member of the TRAF family to be up-regulated in human carcinomas. TRAF4 is required during embryogenesis in key biological processes including the formation of the trachea, the development of the axial skeleton, and the closure of the neural tube. TRAF4 deficiency leads to severe developmental alterations.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66755-1-Ig (TRAF4 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 66755-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human urothelial carcinoma tissue slide using 66755-1-Ig (TRAF4 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66755-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human urothelial carcinoma tissue slide using 66755-1-Ig (TRAF4 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66755-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using 66755-1-Ig (TRAF4 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66755-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66755-1-Ig (TRAF4 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 66755-1-PBS in a different storage buffer formulation.