For Research Use Only

DKK3 Monoclonal antibody

Catalog Number:66758-1-lg 2 Publications




Basic Information	Catalog Number: 66758-1-lg	GenBank Accession Number: BC007660			
	Size: 1200 µ g/ml	GenelD (NCBI): 27122	CloneNo.: 4E6H6		
	Source: Mouse	UNIPROT ID: Q9UBP4	Recommended WB 1:1000-1:6		
	lsotype: lgG2b	Full Name: dickkopf homolog 3 (Xenopus la	IE 1:/00 1:160	IHC 1:150-1:600) IF 1:400-1:1600	
	Immunogen Catalog Number: AG23615	Calculated MW: 38 kDa			
		Observed MW: 40 kDa, 50-55 kDa			
Applications	Tested Applications:	Positive Controls: WB : HEK-293 cells, MCF-7 cells, NCCIT cells			
	IF/ICC, IHC, WB, ELISA				
	Cited Applications: WB, IF	IHC : hu	IHC : human heart tissue, mouse brain tissue		
	Species Specificity: Human, Mouse, Rat, Pig	IF : HEK-293 cells,			
	Cited Species: human, pig				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	DKK3, also named as REIC, belongs to the dickkopf family. It antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6. DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer's dise The adenovirus carrying REIC/Dkk-3 (Ad-REIC) exhibits a potent tumor-specific cell-killing function for various human cancers. It has also become evident that some human cancers are resistant to Ad-REIC-induced apoptosis. The DKK3 is a Glycosylation protein with MW about 55 kDa or 38-43 kDa.				
Notable Publications	Author	Pubmed ID Journal		Application	
	Renwu Hua	36284344 Reprod Biol End	locrinol	WB	
	Michael T H Ng	38374174 Nat Commun		WB,IF	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide an Aliquoting is unnecessary for -20	d 50% glycerol pH 7.3.			

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data





Various lysates were subjected to SDS PAGE Imm followed by western blot with 66758-1-1g (DKK3 emb antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human heart tissue slide using 66758-1-Ig (DKK3 antibody) at dilution of 1:300 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using DKK3 antibody (66758-1-1g, Clone: 4E6H6) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).