For Research Use Only

EEF1A2 Monoclonal antibody, PBS Only



Protein G purification

CloneNo.:

4F3F4

Catalog Number: 66806-1-PBS

Basic Information

Catalog Number: 66806-1-PBS

Size: 1 mg/ml Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG6875

Tested Applications:

WB,Indirect ELISA **Species Specificity:** Human, mouse, rat, pig GenBank Accession Number: **Purification Method:**

BC000432 GeneID (NCBI): 1917 **UNIPROT ID:**

Q05639 Full Name:

eukaryotic translation elongation

factor 1 alpha 2 Calculated MW: 50 kDa

Observed MW: 48-50 kDa

Background Information


Applications

eEF1A2 is one of two isoforms (eEF1A1 and eEF1A2) of eukaryotic elongation factor 1 alpha (eEF1A or eEF1 α). The two isoforms share more than 90% sequence identity and have the similar function that being a protein translation factor involved in protein synthesis. Addition, eEF1A2 plays an important role in cell cycle regulation, heat-shock and the protein synthesis. Addition are provided in the protein synthesis. Addition are provided in provided in protein synthesis. Addition are provided in protein syntresponse, aging, posttranslational modifications, and signal transduction (PMID: 25905039). eEF1A2 is expressed in heart, brain tissue, diaphragm and skeletal muscle while eEF1A1 is expressed ubiquitously (PMID:14588074). And the eEF1A2 protein stimulates the phospholipid signaling and activates the Akt-dependent cell migration and actin $remodeling that \, ultimately \, favors \, tumorigenesis. \, It \, is \, reported \, that \, eEF1A2 \, was \, related \, with \, kinds \, of \, cancers \, such \, respectively. \, The expectation is the expectation of the expectation of$ as ovarian cancer, prostate cancer, pancreatic cancer, breast cancer and lung cancer (PMID:14588074; 24853801; 25744894; 25905039). An 48 kDa band has also been reported(PMID: 24285179).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66806-1-lg (EEF1A2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66806-1-PBS in a different storage buffer formulation.



mouse brain tissue were subjected to SDS PAGE followed by western blot with 66806-1-lg (EEF1A2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66806-1-PBS in a different storage buffer formulation.



rat brain tissue were subjected to SDS PAGE followed by western blot with 66806-1-lg (EEF1A2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66806-1-PBS in a different storage buffer formulation.