For Research Use Only

ESRRB Monoclonal antibody, PBS Only



Purification Method:

Protein A purification

CloneNo.:

3E10C1

Catalog Number: 66818-1-PBS

Basic Information

Catalog Number:

66818-1-PBS BC131517

 Size:
 GeneID (NCBI):

 1 mg/ml
 2103

 Source:
 UNIPROT ID:

 Mouse
 095718

 Isotype:
 Full Name:

IgG1 estrogen-related receptor beta

Immunogen Catalog Number:Calculated MW:AG18626508 aa, 56 kDaObserved MW:

47 kDa

GenBank Accession Number:

Applications

Tested Applications:

WB,Indirect ELISA,IHC,IF Species Specificity:

Human, mouse, rat

Background Information


Estrogen related receptor beta (ESRRB), a NR3 Steroid Receptor, has been shown to affect early placental development. Also it can act as a repressor of transcriptional activity mediated by the glucocorticoid receptor (GR). It binds as a monomer to the extended half-site TNAAGTGCA and as a homodimer to the estrogen response element (ERE), palindromic thyroid hormone response element (TRE(pal)), and SF-1 response element, but not to the glucocorticoid response element (GRE). Estrogen related receptor beta mutations lead to abnormal development of the chorion and defective diploid trophoblast proliferation, and are lethal during midgestation. Expression has been documented in mice in the extra embryonic ectoderm that gives rise to the chorion. ESTs have been isolated from normal human eye, lung, and testis libraries. The ligand for the receptor is PPARgamma coactivator 1 (PGC-1) beta. ESRRB exists various isoforms and molecular weight of each isoform is 48 kDa,55 kDa and 56 kDa.

Storage


Storage:

Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66818-1-lg (ESRRB antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66818-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66818-1-lg (ESRRB antibody) at dilution of 1:40000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66818-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 66818-1-Ig (ESRRB antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66818-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed Caco-2 cells using ESRRB antibody (66818-1-Ig, Clone: 3E10C1) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66818-1-PBS in a different storage buffer formulation.