For Research Use Only

## HDAC4 Monoclonal antibody, PBS Only



Catalog Number:66838-1-PBS


Catalog Number: 66838-1-PBS	GenBank Accession Number: BC039904	Purification Method: Protein A purification
Size: 1mg/ml	GenelD (NCBI): 9759	CloneNo.: 1D2G12
Source: Mouse	UNIPROT ID: P56524	
lsotype: IgG2a	Full Name: histone deacetylase 4	
Immunogen Catalog Number: AG11345	Calculated MW: 972 aa, 106 kDa	
	Observed MW: 140 kDa	
Tested Applications: WB, IHC, Indirect ELISA		
Species Specificity: Human		
deacetylation of lysine residues of deacetylation gives a tag for epig cycle progression and developme	on the N-terminal part of the core histo enetic repression and plays an import ntal events. Histone deacetylases act	nes (H2A, H2B, H3 and H4). Histone ant role in transcriptional regulation, cell via the formation of large multiprotein
Storage: Store at -80°C. Storage Buffer:		
	66838-1-PBS Size: Img/ml Source: Mouse Isotype: IgG2a Immunogen Catalog Number: AG11345 Tested Applications: WB, IHC, Indirect ELISA Species Specificity: Human HDAC4, also named as HDACA, be deacetylation of lysine residues of deacetylation gives a tag for epig cycle progression and developme complexes. HDAC4 is Involved in MEF2A, MEF2C and MEF2D. Storage: Storage: Storage:	66838-1-PBS BC039904   Size: GeneID (NCBI):   Img/ml 9759   Source: UNIPROT ID:   Mouse P56524   Isotype: Full Name:   IgG2a histone deacetylase 4   Immunogen Catalog Number: Calculated MW:   AG11345 972 aa, 106 kDa   Observed MW: 140 kDa   Tested Applications: WB, IHC, Indirect ELISA   Species Specificity: Human   ON HDAC4, also named as HDACA, belongs to the histone deacetylase fam deacetylation of lysine residues on the N-terminal part of the core histor deacetylation gives a tag for epigenetic repression and plays an import cycle progression and developmental events. Histone deacetylases act complexes. HDAC4 is Involved in muscle maturation via its interaction MEF2A, MEF2C and MEF2D.   Storage: Storage:   Storage: Storage:

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 66838-1-Ig (HDAC4 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66838-1-PBS in a different storage buffer formulation.

Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66838-1-1g (HDAC4 antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66838-1-PBS in a different storage buffer formulation.