For Research Use Only

Phospho-RIPK1 (Ser161) Monoclonal antibody, PBS Only

www.ptglab.com

Purification Method:

Protein A purification

CloneNo.:

1B2G1

Catalog Number: 66854-1-PBS

Basic Information

Catalog Number:

66854-1-PBS

Size: 1mg/ml

Source:

Mouse

Isotype: lgG1

NM_003804 GeneID (NCBI):

GenBank Accession Number:

UNIPROT ID: Q13546 Full Name:

receptor (TNFRSF)-interacting serinethreonine kinase 1

Calculated MW: 76 kDa Observed MW: 74 kDa, 45 kDa

Applications

Tested Applications: WB, IHC, ELISA, IF/ICC Species Specificity:

Background Information

RIPK1(Receptor-interacting serine/threonine-protein kinase 1) is primarily involved in mediating TNF-R1-induced cell activation, apoptosis and necroptosis and belongs to a novel class of kinases that function in cell survival and cell death mechanisms(PMID:22685397).It has 2 isoforms produced by alternative splicing.It also can exist as the protein with lower molecular weight about 25kd and 45kd in HAEC and HUVEC(PMID:22685397).


Storage

Storage:


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS Only


Selected Validation Data




Non-treated HEK-293T and Calyculin A treated HEK-293T cells were subjected to SDS PAGE followed by western blot with 66854-1-Ig (Phospho-RIPK1 (Ser161) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control. This data was developed using the same antibody clone with 66854-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 66854-1-lg (Phospho-RIPKI (Ser161) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was untreated (left) or treated with Lambda Protein Phosphatase (λ -PPase, 500U, right) at 37°C for 1 hours before blocking. This data was developed using the same antibody clone with 66854-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 66854-1-Ig (RIPK1 phospho S161 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66854-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 66854-1-lg (RIPK1 phospho S161 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66854-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed U2OS cells using Phospho-RIPK1 (Ser161) antibody (66854-1-Ig, Clone: 1B2G1) at dilution of 1:400 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66854-1-PBS in a different storage buffer formulation.



1X10^6 Calyculin A treated HEK-293T cells were intracellularly stained with 0.13 ug Anti-Human Phospho-RIPK1 (Ser161) (66854-1-lg, Clone:1B2G1) labeled with FlexAble Coralite® Plus 555 Antibody Labeling Kit for Mouse IgG1 (KFA022). Cells were fixed with 49% PFA and permeabilized with 90% MeOH. This data was developed using the same antibody clone with 66854-1-PBS in a different storage buffer formulation.