For Research Use Only

CD8 Monoclonal antibody, PBS Only

Catalog Number: 66868-1-PBS



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

66868-1-PBS

GeneID (NCBI):

BC025715

Protein A purification

Size:

CloneNo.: 1G2B10

1 mg/ml Source:

ENSEMBL Gene ID: ENSG00000153563

Mouse Isotype: lgG1

UNIPROT ID: P01732

Immunogen Catalog Number:

Full Name: CD8a molecule

AG11111

Calculated MW: 235 aa, 26 kDa Observed MW:

32 kDa

Applications

Tested Applications:

WB,IHC,IF,ELISA

Species Specificity: Human

Background Information


CD8 is a transmembrane glycoprotein that is predominantly expressed on the surface of cytotoxic T cells, and can also be found on natural killer cells, cortical thymocytes, and dendritic cells. CD8 serves as a co-receptor for the T cell receptor (TCR). Both CD8 and TCR recognize antigens displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. CD8 plays a role in T cell development and activation of mature T cells. This antibody is raised against 23-183 aa of human CD8a, the alpha chain of CD8 molecule.

Storage


Storage: Store at -80°C.

Storage Buffer: PBS only


Selected Validation Data




MOLT-4 cells were subjected to SDS PAGE followed by western blot with 66868-1-1g (CD8a antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66868-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66868-1-lg (CD8a antibody) at dilution of 1:20000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66868-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66868-1-lg (CD8a antibody) at dilution of 1:20000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66868-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD8a antibody (66868-1-Ig, Clone: 1G2B10) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66868-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD8a antibody (66868-1-lg, Clone: 1G2B10) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66868-1-PBS in a different storage buffer formulation.