

For Research Use Only

TERF2 Monoclonal antibody, PBS Only



Catalog Number: 66893-1-PBS

Basic Information

Catalog Number: 66893-1-PBS	GenBank Accession Number: BC024890	Purification Method: Protein A purification
Size: 1 mg/ml	GeneID (NCBI): 7014	CloneNo.: 5B1E1
Source: Mouse	UNIPROT ID: Q15554	
Isotype: IgG1	Full Name: telomeric repeat binding factor 2	
Immunogen Catalog Number: AG28513	Calculated MW: 56 kDa	
	Observed MW: 60-65.32-35 kDa	

Applications

Tested Applications:
WB, Indirect ELISA, IHC

Species Specificity:
Human, mouse, rat

Background Information

TERF2, also named as TRF2 and Telomeric repeat-binding factor 2, is a 542 amino acid protein, which contains 1 HTH myb-type DNA-binding domain and localizes in the Nucleus. TRF2 binds the telomeric double-stranded 5'-TTAGGG-3' repeat and plays a central role in telomere maintenance and protection against end-to-end fusion of chromosomes. TERF2 is a component of the shelterin complex (telosome) that is involved in the regulation of telomere length and protection. TERF2 together with DCLRE1B/Apollo, is required to control the amount of DNA topoisomerase (TOP1, TOP2A and TOP2B), which is needed for telomere replication during fork passage and prevent aberrant telomere topology. TERF2 recruits TERF2IP/RAP1 to telomeres, thereby participating in to repressing homology-directed repair (HDR), which can affect telomere length. TERF2 exists some isoforms with MV 60 kDa and 32 kDa.

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




MCF-7 cells were subjected to SDS PAGE followed by western blot with 66893-1-Ig (TERF2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66893-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 66893-1-Ig (TERF2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66893-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 66893-1-Ig (TERF2 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66893-1-PBS in a different storage buffer formulation.



HEK-293 cells were subjected to SDS PAGE followed by western blot with 66893-1-Ig (TERF2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66893-1-PBS in a different storage buffer formulation.



Jurkat cells were subjected to SDS PAGE followed by western blot with 66893-1-Ig (TERF2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66893-1-PBS in a different storage buffer formulation.



4T1 cells were subjected to SDS PAGE followed by western blot with 66893-1-Ig (TERF2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66893-1-PBS in a different storage buffer formulation.