

For Research Use Only

UPF1 Monoclonal antibody, PBS Only



Catalog Number: 66898-1-PBS

Featured Product

Basic Information

Catalog Number:

66898-1-PBS

Size:

1 mg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG28320

GenBank Accession Number:

BC039817

GeneID (NCBI):

5976

UNIPROT ID:

Q92900

Full Name:

UPF1 regulator of nonsense transcripts homolog (yeast)

Calculated MW:

123 kDa

Observed MW:

123-130 kDa

Purification Method:

Protein G purification

CloneNo.:

3B6B5

Applications

Tested Applications:

WB, IF, IHC, Indirect ELISA

Species Specificity:

Human, Mouse, Rat

Background Information

Up-Frameshift Suppressor 1 Homolog (UPF1) is the central factor in nonsense-mediated mRNA decay (NMD) and is also directly involved in telomere homeostasis, DNA replication, histone mRNA degradation and staufen-mediated mRNA decay (PMID: 29382845). It is a potential modulator of MALAT1 and that UPF1/MALAT1 pathway could be a therapeutic target for gastric cancer (PMID: 28942451). The molecular mass of UPF1 is 123-130 kDa.

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66898-1-Ig (UPF1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66898-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66898-1-Ig (UPF1 antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66898-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66898-1-Ig (UPF1 antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66898-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66898-1-Ig (UPF1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66898-1-PBS in a different storage buffer formulation.



WB result of UPF1 antibody (66898-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-UPF1 transfected HEK-293 cells. This data was developed using the same antibody clone with 66898-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using UPF1 antibody (66898-1-Ig, Clone: 3B6B5) at dilution of 1:800 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66898-1-PBS in a different storage buffer formulation.