For Research Use Only

TIAR Monoclonal antibody, PBS Only

Catalog Number:66907-1-PBS




Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 66907-1-PBS BC030025 Protein G purification GenelD (NCBI): CloneNo.: Size: 1 mg/ml 7073 1E8D12 Source: UNIPROT ID: Mouse Q01085 Isotype: Full Name: lgG1 TIA1 cytotoxic granule-associated RNA binding protein-like 1 Immunogen Catalog Number: AG11981 Calculated MW: 42 kDa Observed MW: 39 and 41 kDa **Applications Tested Applications:** WB,Indirect ELISA,IHC,IF,FC **Species Specificity:** Human, Mouse, Rat **Background Information** TIAR, also named as TIA 1 related protein, is an RNA-binding protein which is associated with apoptosis. It has been shown that mouse with disrupted TIAR gene show high levels of embryonic lethality. TIAR plays an important role in cytoplasm and nucleus where it can control translation of some specific mRNAs and activate splicing of exons with weak 5-splice sites. 66907-1-Ig detects 39 and 41 kDa bands in SDS-PAGE. Besides, TIAR also has the 28 kDa and 50 kDa proteins for TIAR exists in multiple alternative splicing forms.(PMID: 12533540, 7533298, 22851315, 8176212) Storage: Storage Store at -80°C. Storage Buffer: PBS Only

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66907-1-1g (TIAR antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66907-1-PBS in a different storage buffer formulation.




RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 66907-1-1g (TIAR antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66907-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66907-1-1g (TIAR antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66907-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using 66907-1-Ig (TIAR antibody) at dilution of 1:100. This data was developed using the same antibody clone with 66907-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66907-1-Ig (TIAR antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66907-1-PBS in a different storage buffer formulation.



1X10⁶ MCF-7 cells were intracellularly stained with 0.2 ug Anti-Human TIAR (66907-1-lg, Clone:1E8D12) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 66907-1-PBS in a different storage buffer formulation.