For Research Use Only

ROR1 Monoclonal antibody, PBS Only proteintech

Catalog Number: 66923-1-PBS

Featured Product



Basic Information

Catalog Number:

BC006374

Purification Method:

66923-1-PBS

GeneID (NCBI):

GenBank Accession Number:

Protein A purification

Size: 1 mg/ml

4919

CloneNo.: 1F5D8

1 mg/m Source:

UNIPROT ID: Q01973

Mouse Isotype:

Full Name:

lgG1

receptor tyrosine kinase-like orphan

receptor 1

AG14586

Calculated MW: 937 aa. 104 kDa

Observed MW:

130 kDa

Applications

Tested Applications:

WB, IHC, Indirect ELISA

Immunogen Catalog Number:

Species Specificity:

Human, mouse

Background Information

ROR1 is a member of the RTK family of orphan receptors related to muscle-specific kinase and Trk neurotrophin receptors (PMID: 18546292). ROR1 is primarily expressed by neural crest cells during embryogenesis. High expression of ROR1 is reported in many types of malignancies and is thought to be involved in tumor growth, apoptosis, and epithelial-mesenchymal transition (PMID: 26245996). The human ROR1 gene encodes a 937 amino acid protein with an Ig-like domain, a cysteine-rich domain, a kringle domain, a tyrosine kinase domain, and a proline-rich domain. The calculated molecular weight of ROR1 is 104 kDa, but ROR1 has multiple N-glycosylation sites that generate post-translationally modified ROR1 at 130 kDa (PMID: 24752542).

Storage

Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C


Storage Buffer:

PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66923-1-1g (ROR1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control. This data was developed using the same antibody clone with 66923-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66923-1-lg (ROR1 antibody) at dilution of 1:600 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66923-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66923-1-lg (ROR1 antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66923-1-PBS in a different storage buffer formulation.



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 66923-1-lg (ROR1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66923-1-PBS in a different storage buffer formulation.