## For Research Use Only

## DDX21 Monoclonal antibody, PBS Only



Catalog Number: 66925-1-PBS

**Basic Information** 

Catalog Number: 66925-1-PBS

Size: 1 mg/ml Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG17850

Tested Applications:

WB,Indirect ELISA,IHC,IF,FC Species Specificity:

## **Background Information**

DDX21 protein belongs to DEAD box protein family which is characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD). As a putative RNA helicase, DDX21 unwinds double-stranded RNA, folds single-stranded RNA and is involved in process including ribosomal RNA biogeneis, RNA editing and general transcription. Interaction of DDX21 and c-Jun was reported in ribosomal RNA processing.

GenBank Accession Number:

DEAD (Asp-Glu-Ala-Asp) box

BC008071

9188

GeneID (NCBI):

**UNIPROT ID:** 

polypeptide 21

Calculated MW: 783 aa, 87 kDa Observed MW: 90-95 kDa

Q9NR30 Full Name:

Storage


**Applications** 

Storage: Store at -80°C. Storage Buffer: PBS Only


**Purification Method:** Protein A purification

CloneNo.: 1B9C8


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66925-1-lg (DDX21 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66925-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 66925-1-1g (DDX21 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control. This data was developed using the same antibody clone with 66925-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66925-1-lg (DDX21 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66925-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 66925-1-Ig (DDX21 antibody) at dilution of 1:100 and Coral.ite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66925-1-PBS in a different storage buffer formulation.



1X10^6 HepG2 cells were intracellularly stained with 0.2 ug Anti-Human DDX21 (66925-1-Ig, Clone:1B9C8) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Mouse IgG1 Isotype Control (66360-1-Ig, Clone: T1F8D3F10) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 66925-1-PBS in a different storage buffer