

For Research Use Only

UBAP1 Monoclonal antibody

Catalog Number: 66993-1-Ig



Basic Information

Catalog Number: 66993-1-Ig	GenBank Accession Number: BC020950	Purification Method: Protein G purification
Size: 1900 µg/ml	GeneID (NCBI): 51271	CloneNo.: 2A6A2
Source: Mouse	UNIPROT ID: Q9NZ09	Recommended Dilutions: WB 1:1000-1:6000 IHC 1:50-1:500
Isotype: IgG1	Full Name: ubiquitin associated protein 1	
Immunogen Catalog Number: AG17675	Calculated MW: 502 aa, 55 kDa	
	Observed MW: 55 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, HEK-293 cells, Jurkat cells

IHC : human heart tissue, human lung cancer tissue

Background Information

UBAP1, also named as NAG20, is a 55 kDa protein which contains 2 putative tandem UBA domains at the C terminus, a coiled-coil domain, 2 possible N-glycosylation sites, 5 PKC phosphorylation sites, 7 casein kinase II phosphorylation sites, and 3 N-myristoylation sites. UBAP1 might be a potential effective diagnosis candidate for NPC and decreased expression of UBAP1 protein is a possible point of dysfunction along the pathogenesis pathway for NPC that may contribute to malignant transformation.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66993-1-Ig (UBAP1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human heart tissue slide using 66993-1-Ig (UBAP1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human heart tissue slide using 66993-1-Ig (UBAP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).