

For Research Use Only

TIMELESS Monoclonal antibody, PBS Only

Catalog Number: 67022-1-PBS

Featured Product



Basic Information

Catalog Number: 67022-1-PBS	GenBank Accession Number: BC050557	Purification Method: Protein A purification
Size: 1 mg/ml	GeneID (NCBI): 8914	CloneNo.: 1B6G3
Source: Mouse	UNIPROT ID: Q9UNS1	
Isotype: IgG1	Full Name: timeless homolog (Drosophila)	
Immunogen Catalog Number: AG6283	Calculated MW: 139 kDa	
	Observed MW: 140 kDa	

Applications

Tested Applications:
WB, FC (Intra), Indirect ELISA

Species Specificity:
human, mouse, rat

Background Information

TIMELESS1 (TIM1) gene encodes protein timeless homolog belonging to timeless family. TIM1 is localized in nucleus and regulated by cell cycle. It expresses high levels in S, G2 and M phases, and low in G0 and G1 phases. TIM1 is involved in the circadian rhythm autoregulatory loop and is a crucial component of the circadian core oscillator, which includes the CRY proteins, CLOCK, NPAS2, etc. It is also involved in cell survival after DNA damage or replication stress and reported to play important role in epithelial cell morphogenesis and formation of branching tubules.

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67022-1-Ig (TIMELESS antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control. This data was developed using the same antibody clone with 67022-1-PBS in a different storage buffer formulation.



1×10^6 HEK-293 cells were intracellularly stained with 0.4 ug Anti-Human TIMELESS (67022-1-Ig, Clone:1B6G3) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 67022-1-PBS in a different storage buffer formulation.