## For Research Use Only

## PELP1 Monoclonal antibody, PBS Only



**Purification Method:** 

Protein A purification

CloneNo.:

1B11E4

Catalog Number: 67050-1-PBS

**Basic Information** 

Catalog Number: 67050-1-PBS

Size: 1mg/ml Source: Mouse Isotype:

lgG2b Immunogen Catalog Number:

AG25729

Observed MW: 160 kDa

**Applications** 

**Tested Applications:** Indirect ELISA, IF-P, IHC, WB

Species Specificity:

## **Background Information**

PELP1 was first identified as a 160 kDa protein in a screen for Src homology 2 (SH2) domain-binding proteins. PELP1 is overexpressed in 60-80% of breast tumors and plays important roles in both ER genomic and non-genomic signaling. In vivo, PELP1 subcellular localization is primarily nuclear in normal breast tissue, but it is localized to the cytoplasm in about 40% of invasive breast tumors. In the nucleus, PELP1 interacts with a number of transcription factors. The proto-oncogenic functions of PELP1 involve different cellular processes including epigenetic modifications leading to ER transactivation and breast cancer progression. Furthermore, PELP1 activates kinase cascades in the cytoplasm such as MAPK activation via c-Src and PI3K signaling.

GenBank Accession Number:

proline, glutamate and leucine rich

BC069058

27043

Q8IZL8 Full Name:

protein 1

Calculated MW:

1130 aa, 120 kDa

GeneID (NCBI):

**UNIPROT ID:** 

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

PBS Only


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67050-1-lg (PELP1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 67050-1-lg (PELP1 antibody) at dilution of 1:1000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 67050-1-lg (PELP1 antibody) at dilution of 1:1000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using PELP1 antibody (67050-1-1g, Clone: 1811E4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer formulation.



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human PELP1 (67050-1-lg, Clone:1B11E4) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-lg, Clone: K11B8C4B5) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using PELP1 antibody (67050-1-lg, Clone: 1811E4) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67050-1-PBS in a different storage buffer formulation.