

For Research Use Only

TFAP2A/AP-2 Monoclonal antibody, PBS Only



Catalog Number: 67076-1-PBS

Basic Information

Catalog Number: 67076-1-PBS	GenBank Accession Number: BC017754	Purification Method: Protein A purification
Size: 1 mg/ml	GeneID (NCBI): 7020	CloneNo.: 1B12F11
Source: Mouse	UNIPROT ID: P05549	
Isotype: IgG1	Full Name: transcription factor AP-2 alpha (activating enhancer binding protein 2 alpha)	
Immunogen Catalog Number: AG4112	Calculated MW: 431 aa, 47 kDa	
	Observed MW: 47 kDa	

Applications

Tested Applications:
WB, IHC, ELISA

Species Specificity:
human, mouse, rat

Background Information

The activator protein-2 (AP-2) family of transcription factors comprises five 52-kDa isoforms (AP-2 α , AP-2 β , AP-2 γ , AP-2 δ , and AP-2 ϵ), which share a common structure: a proline/glutamine-rich transactivation domain in the N-terminal region and a helix-span-helix domain in the C-terminal region, which mediates dimerization and site-specific DNA binding. Depending on the cellular context, the AP-2 transcription factors are individually associated either with cell differentiation and development or with cancer progression/regression. [PMID:21966377] TFAP2A (AP-2-alpha) is the only AP-2 protein required for early morphogenesis of the lens vesicle. Together with the CITED2 coactivator, stimulates the PITX2 P1 promoter transcription activation [PMID:11694877]

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



L02 cells were subjected to SDS PAGE followed by western blot with 67076-1-Ig (TFAP2A/AP-2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67076-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 67076-1-Ig (TFAP2A/AP-2 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67076-1-PBS in a different storage buffer formulation.



1X10⁶ A431 cells were intracellularly stained with 0.4 ug Anti-Human TFAP2A/AP-2 (67076-1-Ig, Clone:1B12F11) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 67076-1-PBS in a