For Research Use Only

RUNX1T1 Monoclonal antibody

Catalog Number: 67086-1-Ig



Basic Information

Catalog Number: 67086-1-lg Size: 1400 ug/ml

Source: Mouse Isotype: lgG1 runt-related transcription factor 1;

Immunogen Catalog Number:

AG7893

GenBank Accession Number:

BC005850 GeneID (NCBI): 862

UNIPROT ID: Q06455 **Full Name:**

translocated to, 1 (cyclin D-related) Calculated MW:

Observed MW: 70-75 kDa

68 kDa

Applications

Tested Applications: WB, IHC, ELISA

Species Specificity: human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed

with citrate buffer pH 6.0

Purification Method:

Protein G purification

CloneNo.: 2G2F7

Recommended Dilutions:

WB 1:1000-1:6000 IHC 1:500-1:2000

Positive Controls:

WB: Jurkat cells, HEK-293 cells, pig brain tissue, K-562 cells, rat brain tissue, mouse brain tissue

IHC: mouse brain tissue.

Background Information

RUNX1T1 is a putative zinc finger transcription factor and oncoprotein. In acute myeloid leukemia, especially in the M2 subtype, the t(8;21)(q22;q22) translocation is one of the most frequent karyotypic abnormalities. The translocation produces a chimeric gene made up of the 5'-region of the RUNX1 gene fused to the 3'-region of this $gene.\ Various\ transcript\ of\ the\ fusion\ gene\ has\ been\ reported.\ RUNX1T1\ exists\ some\ isoforms\ with\ MV\ 68,\ 67,64,\ 48$ and 44 kDa. The calcualted molecular weight of RUNX1T1 is 67 kDa, but modified RUNX1T1 is about 70-75 kDa.

Storage


Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data





WB result of RUNX1T1 antibody (67086-1-lg; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-RUNX1T1 transfected HEK-293 cells.

Jurkat cells were subjected to SDS PAGE followed by western blot with 67086-1-1g (RUNX1T1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67086-1-Ig (RUNX1T1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).